

# Drug Information Matrix

| Drug Name  | Substance Detected in Urine   | What to Look For & Physical Symptoms  | Dangers   | Time Detectable in Urine                                  |
|--|---|---|---|---|
| <b>Amphetamines</b><br>·Uppers·Speed·Meth<br>·Whites·Dexies·Crank<br>·Black Beauties       | Methamphetamine, Amphetamine  | Tablets of varying colors, possible chain smoking, long periods without rest or sleep.<br>Loss of appetite, anxiety, irritability, rapid speech, tremors, mood elevation.   | Disorientation, severe depression, paranoia; possible hallucinations, increase in blood pressure, fatigue.  | 1 to 2 days   |
| <b>Cocaine</b><br>·Coke·Crack·Snow·C<br>·Flake·White Candy<br>·Free Base·Blow              | Benzoylcegonine   | Glassine envelopes, razor, small spoons; odorless, bitter white crystalline powder, granular rocks. Short-lived euphoria changing to depression, irritability, nervousness, tightness of muscles.   | Shallow breathing, fever, anxiety, tremors, possible death from convulsions or respiratory arrest   | 1 to 3 days   |
| <b>Marijuana</b><br>·Dope·Pot·Reefer<br>·Joint·Grass·Toot·Blow                             | 11-Nor-delta-9-tetrahydrocannabinol-9-carboxylic acid                               | Plastic baggies, rolling paper, "roach" clips, odor of burnt rope. Altered perception, dilated pupils, lack of concentration and coordination, craving for sweets, increased appetite, laughter   | Psychological dependence, increased heart rate, impaired short-term memory, anxiety, lung damage, possible psychosis with chronic use.                                  | Occasional Use: 1 to 3 days<br>Chronic Use: Up to 30 days |
| <b>Opiates</b><br>·Heroin (smack, horse, junk)·Morphine ("M")<br>·Codeine (school boy)     | Codeine, Morphine   | Glassine envelopes, needles and syringes, caps or spoons, tourniquets, needle marks on body. Insensitivity to pain, euphoria, sedation, nausea, vomiting, itchiness, watery eyes, running nose  | Lethargy, weight loss, hepatitis, slow and shallow breathing, possible death.   | 1 to 3 days   |
| <b>Phencyclidine</b><br>·Angel Dust·Devil Stick<br>·PCP·Dummy Dust                         | Phencyclidine   | Liquid capsules. White or brown powder can be put on paper stamps, sugar cubes, cigarettes or joints. May be injected. Increased pulse and heart rate, blood pressure and temperature. Mood and perception alteration possible, paranoia, panic, anxiety, nausea, tremors, suicidal urge. | Unpredictable behavior, flash backs, possible emotional instability and psychosis, hallucination.   | Occasional Use: 1 to 5 days<br>Chronic Use: Up to 30 days |
| <b>Barbiturates</b><br>·Downers·Dolls·Reds<br>·Tunia·Rainbows·Yellows<br>·Blues·Goof Balls | Secobarbital<br>Phenobarbital<br>Mephobarbital<br>Butobarbital<br>Amobarbital       | Capsules of varying colors, longer periods of rest or sleep, dizziness, cold and clammy skin.<br>Depression, decreased alertness and muscle control, intoxication and slurred speech, drowsiness  | Rigidity and painful muscle contraction, emotional instability, possible overdose and death, especially when mixed with alcohol.  | Pentobarbital: 1 to 3 days<br>Phenobarbital: 1 to 3 weeks |
| <b>Benzodiazepines</b><br>·Downers   | Diazepam, Chlordiazepoxide, Medazepam, Oxazepam, Nordiazepam, Flurazepam, Temazepam | Oral/injection.<br>Slurred speech, disorientation, drunken behavior without odor of alcohol.  | Shallow respiration, cold and clammy skin, dilated pupils, weak and rapid pulse, coma, possible death.  | 1 to 14 days.   |
| <b>Methadone</b><br>·Done·Dolophine<br>·Methadose  | Methadone   | Tablets, liquid, injection.<br>Euphoria, drowsiness, respiratory depression, constricted pupils, nausea.  | Slow, shallow breathing, clammy skin, convulsions, coma, possible death.  | 1 to 3 days   |
| <b>Methaqualone</b>  | Methaqualone, Quaalude, Meguin  | Tablets.<br>Slow heart rate and breathing, lowered blood pressure. Sleepiness, feeling of well being, loss of coordination, dizziness, impaired perception, confusion, later hangover.  | Anxiety, insomnia.  | 1 to 7 days   |
| <b>Propoxyphene</b>  | NorPropoxyphene   | Tablets, capsules.<br>Dizziness, drowsiness, headache, euphoria, dysphoria, asthenia.   | Skin rash and other allergic reactions occur occasionally and may be accompanied by drug fever and mucosal lesion, stupor or coma; convulsions, respiratory depression. | 1 to 3 days.  |

\*These times are estimates, the time determined in urine is affected by various factors including amount and frequency of usage, weight of the individual, other medications taken, etc. You should contact a physician knowledgeable in substance abuse if you have any questions.